

OR-5. NEW APPROACH TO UNSYMMETRICAL 1,3- AND 1,4-DIAZATRIPHENYLENE DERIVATIVES THROUGH INTRAMOLECULAR OXIDATIVE CYCLODEHYDROGENATION

I. M. Pakhomov², E. V. Verbitskiy^{1,2}, Yu. A. Kvashnin¹, A. R. Tameev³,
G. L. Rusinov^{1,2}, O. N. Chupakhin^{1,2}, V. N. Charushin^{1,2}

¹ I. Ya. Postovsky Institute of Organic Synthesis UB RAS,
S. Kovalevskoy/Akademicheskaya St., 20/22, Yekaterinburg, 620990, Russia


² Ural Federal University of the first President of Russia B. N. Yeltsin,
Mira St., 19, Yekaterinburg, 620002, Russia

³ Frumkin Institute of Physical Chemistry and Electrochemistry RAS,
Leninsky prospekt, 31, bld.4, Moscow, 119071, Russia

E-mail: Verbitsky@ios.uran.ru

Polycyclic heteroaromatic compounds have received much attention as a fundamental framework in organic functional materials because their electrochemical and photochemical properties are useful in organic electronics and luminescent materials. Among them, 1,3- and 1,4-diazatriphenylene derivatives frequently appeared in molecular unit in the preparation of extended π -conjugated systems with characteristic optical and electrochemical properties [1].

Herein, we describe the synthesis of an unsymmetrical substituted dibenzo[*f, h*]quinazolines ($X=N$, $Y=CH$, $A=B=C$ – not present) and dibenzo[*f, h*]furazano[3,4-*b*]quinoxalines ($X=A=C=N$, $Y=CH$, $B=O$) by the sequence of Suzuki cross-coupling reaction and intramolecular oxidative cyclodehydrogenation (S_N^H -reaction) [2]. The optimization of reaction conditions for each stage were produced.



The electrochemical and photophysical properties for several polycycles have been measured. In particular, dibenzo[*f, h*]furazano[3,4-*b*]quinoxalines possess hole mobility of the order of $10^{-4} \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$ that make them perspective for fabrication of hole transport layer in perovskite solar cells, because their hole mobility values are close to the reference molecule spiro-OMeTAD and HOMO levels match well with that of MAPbI₃.

References

1. Verbitskiy E. V., Rusinov G. L., Charushin V. N. Diazatriphenylenes and their thiophene analogues: synthesis and applications // *Arkivoc*. 2017. Vol. 2017, № 1. P. 356–401.
2. Charushin V. N., Chupakhin O. N. Metal Free C-H functionalization of Aromatics Nucleophilic Displacement of Hydrogen. Springer : Switzerland, 2014. P. 238.

The research was financially supported by the Russian Science Foundation (Project № 16-13-10435). NMR spectra and elemental analyses were covered by Program of the Ural Branch of the Russian Academy of Sciences (Project № 18-3-3-12).